



198B	Floyd loam, 1 to 4 percent slopes	4.27	5.6%		llw		89	76										
809B	Bertram fine sandy loam, 2 to 5 percent slopes	4.14	5.4%		IVs		39	29										
110B	Lamont fine sandy loam, 2 to 5 percent slopes	3.92	5.1%		llle		52	47										
175B	Dickinson fine sandy loam, 2 to 5 percent slopes	2.77	3.6%		llle		50	55										
407B	Schley loam, 1 to 4 percent slopes	2.28	3.0%		llw		81	73										
626	Hayfield loam, 0 to 2 percent slopes, rarely flooded	2.24	2.9%		lls		53	67										
409B	Dickinson fine sandy loam, loam substratum, 2 to 5 percent slopes	2.16	2.8%		llle		75	56										
177B	Saude loam, 2 to 5 percent slopes	1.75	2.3%		lls		55	60										
84	Clyde silty clay loam, 0 to 3 percent slopes	1.37	1.8%		llw		88	77	4.4	7	4.2	3	193.5	18	3.3	76.5		
1486	Spillville-Sigglekov complex, 0 to 2 % slopes, frequently flooded	0.10	0.1%		Vlllw		5											
Weighted Average							61.6	*-	0.1	0.1	0.1	0.1	3.5	0.3	0.1	1.4		

**IA has updated the CSR values for each county to CSR2.

*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.