





84	Clyde silty clay loam, 0 to 3 percent slopes	4.64	6.4%		llw		88	77	4.4	7	4.2	3	193.5	18	3.3	76.5
407B	Schley loam, 1 to 4 percent slopes	3.16	4.4%		llw		81	73								
391B	Clyde-Floyd complex, 1 to 4 percent slopes	2.98	4.1%		llw		87	73								
41C	Sparta loamy fine sand, 5 to 9 percent slopes	2.03	2.8%		IVs	lle	34	25								
83C	Kenyon loam, 5 to 9 percent slopes	1.73	2.4%		llle		85	72		10.6	6.4		188		3.7	90.5
41B	Sparta loamy fine sand, 2 to 5 percent slopes	0.62	0.9%		IVs	lle	39	40								
284C2	Flagler sandy loam, 5 to 9 percent slopes, moderately eroded	0.33	0.5%		llle		43	15								
393C	Sparta loamy fine sand, loamy substratum, 5 to 9 percent slopes	0.30	0.4%		IVs	lle	33	27								
175B	Dickinson fine sandy loam, 2 to 5 percent slopes	0.14	0.2%		llle		50	55								
<b>Weighted Average</b>							<b>78.3</b>	<b>*-</b>	<b>0.3</b>	<b>0.7</b>	<b>0.4</b>	<b>0.2</b>	<b>17</b>	<b>1.2</b>	<b>0.3</b>	<b>7.1</b>

\*\*IA has updated the CSR values for each county to CSR2.

\*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.