





391B	Clyde-Floyd complex, 1 to 4 percent slopes	10.06	4.6%		llw	87	78										
83B	Kenyon loam, 2 to 5 percent slopes	5.99	2.7%		lle	90	83		11	6.6		210		3.9	91		
177	Saude loam, 0 to 2 percent slopes	2.92	1.3%		lls	60	63										
174B	Bolan loam, 2 to 5 percent slopes	2.36	1.1%		lls	64	70										
175B	Dickinson fine sandy loam, 2 to 5 percent slopes	1.35	0.6%		llle	50	55										
399	Readlyn silt loam, 1 to 3 percent slopes	0.51	0.2%		lw	91	88										
177B	Saude loam, 2 to 5 percent slopes	0.17	0.1%		lls	55	58										
214B	Rockton loam, 20 to 30 inches to limestone, till plain, 2 to 5 percent slopes	0.09	0.0%		lle	47	56										
<b>Weighted Average</b>						<b>72</b>	<b>68.9</b>	<b>0.6</b>	<b>1.3</b>	<b>0.8</b>	<b>0.4</b>	<b>33.5</b>	<b>2.6</b>	<b>0.6</b>	<b>13.5</b>		

\*\*IA has updated the CSR values for each county to CSR2.

\*c: Using Capabilities Class Dominant Condition Aggregation Method  
Soils data provided by USDA and NRCS.